

Kay Dee 20 Block with Bovatec^{®*}



TYPE C MEDICATED

For increased weight gain in pasture cattle,
slaughter, stocker, feeder cattle and dairy and beef replacement heifers.

ACTIVE DRUG INGREDIENT

Lasalocid 300 grams/ton

GUARANTEED ANALYSIS

Crude Protein, not less than 20.0%
(this includes not more than 6.8% crude protein from
non-protein nitrogen)
Crude Fat, not less than 2.0%
Crude Fiber, not more than 4.8%
Calcium (Ca), not less than 1.3%
Calcium (Ca), not more than 1.8%
Phosphorus (P), not less than 1.0%
Salt (NaCl), not less than 18.0%
Salt (NaCl), not more than 21.0%
Magnesium (Mg), not less than 0.01%
Potassium (K), not less than 2.0%
Sulfur (S), not less than 0.035%
Cobalt (Cop), not less than 5 ppm
Copper (Cu), not less than 15 ppm
Iodine (I), not less than 10 ppm
Iron (Fe), not less than 100 ppm
Manganese (Mn), not less than 80 ppm
Zinc (Zn), not less than 500 ppm
Vitamin A, not less than 80,000 IU/lb
Vitamin D₃, not less than 12,000 IU/lb

INGREDIENTS

Grain Products, Plant Protein Products, Salt, Molasses Products, Processed Grain By-Products, Roughage Products, Dicalcium Phosphate, Potassium Chloride, Urea, Sodium Bentonite, Calcium Carbonate, Sodium Sulfate, Vegetable Oil, Zinc Oxide, Iron Oxide, Magnesium Oxide, Manganous Oxide, Copper Oxide, Ethylenediamine Dihydriodide, Cobalt Carbonate, Vitamin A Supplement, Vitamin D Supplement.



Manufactured by

Kay Dee Feed Company
1919 Grand Ave • Sioux City, Iowa 51106

NET WEIGHT 331/3 lbs. (15.13 kg)

FEEDING DIRECTIONS TYPE C MEDICATED FEED

Medicated blocks, as with all self-fed cattle products, must be provided to animals which are supplied with adequate roughage and/or forage at all times.

Self-feed *Kay Dee 20 Block with Bovatec[®]* as the sole source of salt to cattle. Locate blocks near water supply and in cattle loafing areas permitting at least one block for every 10 head of cattle. Maintain adequate *Kay Dee 20 Block with Bovatec[®]* supply at all times to assure continuous intake by adding blocks when each block is about half consumed.

It is essential that cattle fed medicated feed blocks have been adapted to non-medicated block consumption for at least two weeks prior to medicated block feeding. Cattle that are not adapted to block supplementation may not consume the product at the proper level or they may over-consume the product due to inadequated prior supplementation.

Adequate *Kay Dee 20 Block with Bovatec[®]* consumption is required at all times to assure effective drug activity. It is essential that good block husbandry practices be followed at all times. Block feeding techniques include but are not limited to variety of practices. Increased consumption may often be obtained by moving feeding stations into shaded or loafing areas, closer to water source, increasing blocks available to the animals or combinations of these practices. Decreased consumption may often be obtained by the reverse of the above practices.

Kay Dee 20 Block with Bovatec[®] should always be fed in protected feeders.

Adequate pasture/roughage should be available to assure cattle will consume not less than 7 ounces, nor more than 1 pound 5 ounces per head per day.

CAUTION: The safety of Lasalocid for use in unapproved species has not been established. DO NOT allow horses or other equines access to Lasalocid as ingestion may be fatal.

*Bovatec[®] is a registered trademark for Alpha Animal Health's brand of Lasalocid.

NOTICE: This product contains no prohibited animal by-products in accordance with FDA Regulation 21 CFR 589.2000. This manufacturing facility does not handle, store or use any prohibited animal by-products.

Product No. 280-080401

Lot No.
4655



7 60932 00280 9