



20% BLOCK WITH BOVATEC®

**FOR INCREASED WEIGHT GAIN IN PASTURE CATTLE,
SLAUGHTER, STOCKER, FEEDER CATTLE, AND BEEF REPLACEMENT HEIFERS.
A PROTEIN SUPPLEMENT FOR CATTLE ON PASTURE OR IN CONFINEMENT.**

ACTIVE DRUG INGREDIENT

Lasalocid 300 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than 20.00%
(this includes not more than 6.80% crude protein from non-protein nitrogen)
Crude Fat, not less than 2.00%
Crude Fiber, not more than 4.80%
Calcium (Ca), not less than 1.30%
Calcium (Ca), not more than 1.80%
Phosphorus (P), not less than 1.00%
Salt (NaCl), not less than 18.00%
Salt (NaCl), not more than 21.00%
Magnesium (Mg), not less than 0.01%
Potassium (K), not less than 2.00%
Sulfur (S), not less than 0.035%
Cobalt (Co), not less than 5 ppm
Copper (Cu), not less than 15 ppm
Iodine (I), not less than 10 ppm
Iron (Fe), not less than 100 ppm
Manganese (Mn), not less than 80 ppm
Zinc (Zn), not less than 500 ppm
Vitamin A, not less than 80,000 IU/lb
Vitamin D₃, not less than 12,000 IU/lb

FEEDING DIRECTIONS

Medicated blocks, as with all self-fed cattle products, must be provided to animals which are supplied with adequate roughage and/or forage at all times.

Self-feed Kay Dee 20 Block with Bovatec® as the sole source of salt to cattle. Locate blocks near water supply and in cattle loafing areas permitting at least one block for every 10 head of cattle. Maintain adequate Kay Dee 20 Block with Bovatec® supply at all times to assure continuous intake by adding blocks when each block is about half consumed. It is essential that cattle fed medicated feed blocks have been adapted to non-medicated block consumption for at least two weeks prior to medicated block feeding. Cattle that are not adapted to block supplementation may not consume the product at the proper level or they may over-consume the product due to inadequate prior supplementation.

INGREDIENTS

Grain Products, Plant Protein Products, Salt, Molasses Products, Processed Grain By-Products, Roughage Products, Dicalcium Phosphate, Potassium Chloride, Urea, Sodium Bentonite, Calcium Carbonate, Sodium Sulfate, Vegetable Oil, Zinc Oxide, Iron Oxide, Magnesium Oxide, Manganous Oxide, Copper Oxide, Ethylenediamine Dihydriodide, Cobalt Carbonate, Vitamin A Supplement, Vitamin D₃ Supplement.

Adequate Kay Dee 20 Block with Bovatec® consumption is required at all times to assure effective drug activity. It is essential that good block husbandry practices be followed at all times. Block feeding techniques include but are not limited to variety of practices. Increased consumption may often be obtained by moving feeding stations into shaded or loafing areas, closer to water source, increasing blocks available to the animals or combinations of these practices. Decreased consumption may often be obtained by the reverse of the above practices.

Kay Dee 20 Block with Bovatec® should always be fed in protected feeders.

Adequate pasture/roughage should be available to assure cattle will consume not less than 7 ounces, nor more than 1 pound 5 ounces per head per day.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equine access to Type A articles or Type B feeds containing lasalocid, as ingestion may be fatal. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Safety of lasalocid for use in unapproved species has not been established. If overconsumption occurs, limit access to product for one week or until recommended consumption is met.

NOTICE: This product contains no prohibited animal by-products in accordance with FDA Regulation 21 CFR 589.2000. This manufacturing facility does not handle, store or use any prohibited animal by-products.

*Bovatec® is a registered trademark for Alpha Animal Health's brand of Lasalocid.

SPECIFICATIONS:	USE FOR:	FEATURES & BENEFITS:
<ul style="list-style-type: none"> • 33 1/3 lb. block • Feed free choice • Adapt animals to non-medicated blocks before feeding medicated blocks 	<ul style="list-style-type: none"> • For increased weight gain in pasture cattle, slaughter, stocker, feeder cattle, and beef replacement heifers • Promotes optimum condition before calving 	<ul style="list-style-type: none"> • Contains the ionophore Bovatec® to improve feed efficiency and rate of gain • 20% protein, 6.8% NPN • Promotes more efficient use of protein • Ca:P ratio of 1.5:1 • Contains minerals, trace minerals and vitamins
PRODUCT 280		